

$$7.3 \quad f(x) = \cos(x)$$

$$f'(x) = (\cos(x))' = -\sin(x)$$

$$f''(x) = (-\sin(x))' = -\cos(x)$$

$$f^{(3)}(x) = (-\cos(x))' = \sin(x)$$

$$f^{(4)}(x) = (\cos(x))' = \cos(x) = f(x)$$

Plus généralement, on obtient :

$$f^{(k)}(x) = \begin{cases} -\sin(x) & \text{si } k \equiv 1 \pmod{4} \\ -\cos(x) & \text{si } k \equiv 2 \pmod{4} \\ \sin(x) & \text{si } k \equiv 3 \pmod{4} \\ \cos(x) & \text{si } k \equiv 0 \pmod{4} \end{cases}$$

Il en résulte :

$$f^{(k)}(0) = \begin{cases} 0 & \text{si } k \equiv 1 \pmod{4} \\ -1 & \text{si } k \equiv 2 \pmod{4} \\ 0 & \text{si } k \equiv 3 \pmod{4} \\ 1 & \text{si } k \equiv 0 \pmod{4} \end{cases}$$

$$\begin{aligned} 1) \quad P_1(x) &= f(a) + f'(a)(x-a) \\ &= 1 + 0(x-0) \\ &= 1 \end{aligned}$$

$$\begin{aligned} 2) \quad P_2(x) &= f(a) + f'(a)(x-a) + \frac{f''(a)}{2!}(x-a)^2 \\ &= 1 + 0(x-0) + \frac{-1}{2!}(x-0)^2 \\ &= 1 - \frac{x^2}{2!} \end{aligned}$$

$$\begin{aligned} 3) \quad P_3(x) &= f(a) + f'(a)(x-a) + \frac{f''(a)}{2!}(x-a)^2 + \frac{f^{(3)}(a)}{3!}(x-a)^3 \\ &= 1 + 0(x-0) + \frac{-1}{2!}(x-0)^2 + \frac{0}{3!}(x-0)^3 \\ &= 1 - \frac{x^2}{2!} \end{aligned}$$

$$\begin{aligned} 4) \quad P_4(x) &= f(a) + f'(a)(x-a) + \frac{f''(a)}{2!}(x-a)^2 + \frac{f^{(3)}(a)}{3!}(x-a)^3 + \frac{f^{(4)}(a)}{4!}(x-a)^4 \\ &= 1 + 0(x-0) + \frac{-1}{2!}(x-0)^2 + \frac{0}{3!}(x-0)^3 + \frac{1}{4!}(x-0)^4 \\ &= 1 - \frac{x^2}{2!} + \frac{x^4}{4!} \end{aligned}$$

$$\begin{aligned} 5) \quad P_5(x) &= f(a) + f'(a)(x-a) + \frac{f''(a)}{2!}(x-a)^2 + \frac{f^{(3)}(a)}{3!}(x-a)^3 \\ &\quad + \frac{f^{(4)}(a)}{4!}(x-a)^4 + \frac{f^{(5)}(a)}{5!}(x-a)^5 \\ &= 1 + 0(x-0) + \frac{-1}{2!}(x-0)^2 + \frac{0}{3!}(x-0)^3 \\ &\quad + \frac{1}{4!}(x-0)^4 + \frac{0}{5!}(x-0)^5 \\ &= 1 - \frac{x^2}{2!} + \frac{x^4}{4!} \end{aligned}$$

$$\begin{aligned}
6) \quad P_6(x) &= f(a) + f'(a)(x-a) + \frac{f''(a)}{2!}(x-a)^2 + \frac{f^{(3)}(a)}{3!}(x-a)^3 \\
&\quad + \frac{f^{(4)}(a)}{4!}(x-a)^4 + \frac{f^{(5)}(a)}{5!}(x-a)^5 + \frac{f^{(6)}(a)}{6!}(x-a)^6 \\
&= 1 + 0(x-0) + \frac{-1}{2!}(x-0)^2 + \frac{0}{3!}(x-0)^3 \\
&\quad + \frac{1}{4!}(x-0)^4 + \frac{0}{5!}(x-0)^5 + \frac{-1}{6!}(x-0)^6 \\
&= 1 - \frac{x^2}{2!} + \frac{x^4}{4!} - \frac{x^6}{6!}
\end{aligned}$$

7) On généralise facilement le résultat :

$$P_{2n}(x) = 1 - \frac{x^2}{2!} + \frac{x^4}{4!} - \frac{x^6}{6!} + \frac{x^8}{8!} - \frac{x^{10}}{10!} + \frac{x^{12}}{12!} - \frac{x^{14}}{14!} + \dots + (-1)^n \frac{x^{2n}}{(2n)!}$$