

10.15 1) $f'(x) = \sin(2x)$

$$\begin{aligned} f(x) &= \int \sin(2x) dx = \int \sin(2x) \cdot 2 \cdot \frac{1}{2} dx = \frac{1}{2} \int \sin(2x) \cdot 2 dx \\ &= \frac{1}{2} (-\cos(2x)) = -\frac{1}{2} \cos(2x) \end{aligned}$$

$$g(x) = x$$

$$g'(x) = 1$$

$$\begin{aligned} \int x \sin(2x) dx &= -\frac{1}{2} \cos(2x) \cdot x - \int -\frac{1}{2} \cos(2x) \cdot 1 dx \\ &= -\frac{1}{2} x \cos(2x) + \frac{1}{2} \int \cos(2x) dx \\ &= -\frac{1}{2} x \cos(2x) + \frac{1}{2} \int \cos(2x) \cdot 2 \cdot \frac{1}{2} dx \\ &= -\frac{1}{2} x \cos(2x) + \frac{1}{2} \cdot \frac{1}{2} \int \cos(2x) \cdot 2 dx \\ &= -\frac{1}{2} x \cos(2x) + \frac{1}{4} \sin(2x) + c \end{aligned}$$

2) $f'(x) = e^x \quad f(x) = e^x$

$$g(x) = x \quad g'(x) = 1$$

$$\int x e^x dx = x e^x - \int e^x dx = x e^x - e^x = (x-1) e^x + c$$

3) $f'(x) = e^{3x}$

$$f(x) = \int e^{3x} dx = \int e^{3x} \cdot 3 \cdot \frac{1}{3} dx = \frac{1}{3} \int e^{3x} \cdot 3 dx = \frac{1}{3} e^{3x}$$

$$g(x) = 3x^2 \quad g'(x) = 6x$$

$$\int 3x^2 e^{3x} dx = \frac{1}{3} e^{3x} \cdot 3x^2 - \int \frac{1}{3} e^{3x} \cdot 6x dx = x^2 e^{3x} - \int 2x e^{3x} dx$$

Pour calculer $\int 2x e^{3x} dx$, on fait à nouveau une intégration par parties :

$$f'(x) = e^{3x} \quad f(x) = \frac{1}{3} e^{3x}$$

$$g(x) = 2x \quad g'(x) = 2$$

$$\begin{aligned} \int 2x e^{3x} dx &= \frac{1}{3} e^{3x} \cdot 2x - \int \frac{1}{3} e^{3x} \cdot 2 dx = \frac{2}{3} x e^{3x} - \frac{2}{3} \int e^{3x} dx \\ &= \frac{2}{3} x e^{3x} - \frac{2}{3} \cdot \frac{1}{3} e^{3x} = \frac{2}{3} x e^{3x} - \frac{2}{9} e^{3x} \end{aligned}$$

$$\text{Finalement, } \int 3x^2 e^{3x} dx = x^2 e^{3x} - (\frac{2}{3} x e^{3x} - \frac{2}{9} e^{3x})$$

$$= x^2 e^{3x} - \frac{2}{3} x e^{3x} + \frac{2}{9} e^{3x}$$

$$= \frac{1}{9} (9x^2 - 6x + 2) e^{3x} + c$$

$$4) \quad f'(x) = \cos(x) \quad f(x) = \sin(x) \\ g(x) = x^2 + 1 \quad g'(x) = 2x$$

$$\int (x^2 + 1) \cos(x) dx = (x^2 + 1) \sin(x) - \int 2x \sin(x) dx$$

Pour calculer $\int 2x \sin(x) dx$, on procède à une nouvelle intégration par parties :

$$f'(x) = \sin(x) \quad f(x) = -\cos(x) \\ g(x) = 2x \quad g'(x) = 2$$

$$\int 2x \sin(x) dx = -2x \cos(x) - \int -2 \cos(x) dx = -2x \cos(x) + 2 \sin(x)$$

$$\text{Donc } \int (x^2 + 1) \cos(x) dx = (x^2 + 1) \sin(x) - (-2x \cos(x) + 2 \sin(x)) \\ = (x^2 - 1) \sin(x) + 2x \cos(x) + c$$

$$5) \quad f'(x) = 1 \quad f(x) = x \\ g(x) = \ln(x) \quad g'(x) = \frac{1}{x}$$

$$\int \ln(x) dx = x \ln(x) - \int x \cdot \frac{1}{x} dx = x \ln(x) - \int 1 dx = x \ln(x) - x \\ = x (\ln(x) - 1) + c$$

$$6) \quad f'(x) = \sqrt{x+1}$$

$$f(x) = \int \sqrt{x+1} dx = \int (x+1)^{\frac{1}{2}} dx = \frac{1}{\frac{3}{2}} (x+1)^{\frac{3}{2}} = \frac{2}{3} \sqrt{(x+1)^3} \\ = \frac{2}{3} (x+1) \sqrt{x+1}$$

$$g(x) = x \quad g'(x) = 1$$

$$\int x \sqrt{x+1} dx = \frac{2}{3} x (x+1) \sqrt{x+1} - \int \frac{2}{3} (x+1) \sqrt{x+1} dx \\ = \frac{2}{3} x (x+1) \sqrt{x+1} - \frac{2}{3} \int (x+1)^{\frac{3}{2}} dx \\ = \frac{2}{3} x (x+1) \sqrt{x+1} - \frac{2}{3} \cdot \frac{1}{\frac{5}{2}} (x+1)^{\frac{5}{2}} \\ = \frac{2}{3} x (x+1) \sqrt{x+1} - \frac{4}{15} (x+1)^2 \sqrt{x+1} \\ = \frac{2}{15} (x+1) \sqrt{x+1} (5x - 2(x+1)) \\ = \frac{2}{15} (3x - 2) (x+1) \sqrt{x+1} + c$$

$$7) \quad f'(x) = 1$$

$$g(x) = \arcsin(x) \quad g'(x) = \frac{1}{\sqrt{1-x^2}}$$

$$\begin{aligned} \int \arcsin(x) dx &= x \arcsin(x) - \int \frac{1}{\sqrt{1-x^2}} \cdot x dx \\ &= x \arcsin(x) - \left(-\frac{1}{2}\right) \int (1-x^2)^{-\frac{1}{2}} \cdot (-2x) dx \\ &= x \arcsin(x) + \frac{1}{2} \cdot \frac{1}{\frac{1}{2}} (1-x^2)^{\frac{1}{2}} \\ &= x \arcsin(x) + \sqrt{1-x^2} + c \end{aligned}$$

$$8) \quad f'(x) = \cos(x) \quad f(x) = \sin(x)$$

$$g(x) = e^x \quad g'(x) = e^x$$

$$\int e^x \cos(x) dx = e^x \sin(x) - \int e^x \sin(x) dx$$

Pour calculer $\int e^x \sin(x) dx$, on recourt derechef à une intégration par parties :

$$f'(x) = \sin(x) \quad f(x) = -\cos(x)$$

$$g(x) = e^x \quad g'(x) = e^x$$

$$\int e^x \sin(x) dx = -e^x \cos(x) - \int -e^x \cos(x) dx = -e^x \cos(x) + \int e^x \cos(x) dx$$

$$\begin{aligned} \text{Ainsi } \int e^x \cos(x) dx &= e^x \sin(x) - \left(-e^x \cos(x) + \int e^x \cos(x) dx \right) \\ &= e^x (\sin(x) + \cos(x)) - \int e^x \cos(x) dx \end{aligned}$$

$$\text{Il en résulte } 2 \int e^x \cos(x) dx = e^x (\sin(x) + \cos(x))$$

$$\text{d'où finalement } \int e^x \cos(x) dx = \frac{1}{2} e^x (\sin(x) + \cos(x)) + c$$

$$9) \quad f'(x) = x \quad f(x) = \frac{1}{2} x^2$$

$$g(x) = \ln(x) \quad g'(x) = \frac{1}{x}$$

$$\begin{aligned} \int x \ln(x) dx &= \frac{1}{2} x^2 \ln(x) - \int \frac{1}{2} x^2 \cdot \frac{1}{x} dx = \frac{1}{2} x^2 \ln(x) - \frac{1}{2} \int x dx \\ &= \frac{1}{2} x^2 \ln(x) - \frac{1}{4} x^2 = \frac{1}{4} x^2 (2 \ln(x) - 1) + c \end{aligned}$$

$$10) \quad f'(x) = e^{-x} \quad f(x) = -e^{-x}$$
$$g(x) = x^2 \quad g'(x) = 2x$$

$$\int x^2 e^{-x} dx = -x^2 e^{-x} - \int -2x e^{-x} dx = -x^2 e^{-x} + 2 \int x e^{-x} dx$$

Pour calculer $\int x e^{-x} dx$, on procède encore par intégration par parties :

$$f'(x) = e^{-x} \quad f(x) = -e^{-x}$$

$$g(x) = x \quad g'(x) = 1$$

$$\int x e^{-x} dx = -x e^{-x} - \int -e^{-x} dx = -x e^{-x} - e^{-x}$$

$$\text{D'où } \int x^2 e^{-x} dx = -x^2 e^{-x} + 2(-x e^{-x} - e^{-x}) = -(x^2 + 2x + 2)e^{-x} + c$$